

EDGE RINGS SATISFYING SERRE'S CONDITION (R_1)

TAKAYUKI HIBI AND LUKAS KATTHÄN

ABSTRACT. A combinatorial criterion for the edge ring of a finite connected graph to satisfy Serre's condition (R_1) is studied.

INTRODUCTION

The study on edge polytopes and edge rings of finite connected graphs has been achieved from the viewpoints of both combinatorics and computational commutative algebra ([3], [4]). Especially, a combinatorial characterization for the edge ring to be normal is obtained by both [3] and [6] independently. It follows immediately from [2, Theorem 6.4.2] that a normal edge ring is Cohen–Macaulay. However, in general it seems unclear when the edge ring is Cohen–Macaulay. Recall that a noetherian ring is normal if and only if it satisfies Serre's conditions (R_1) and (S_2) . Thus in particular an edge ring satisfying Serre's condition (R_1) is normal if and only if it is Cohen–Macaulay. In the present paper the problem when a given edge ring satisfies Serre's condition (R_1) is investigated.

1. EDGE RINGS AND EDGE POLYTOPES OF FINITE CONNECTED GRAPHS

First, we recall from [3] what edge rings and edge polytopes of finite connected graphs are. Let G be a finite connected graph on the vertex set $[d] = \{1, \dots, d\}$ with $E(G) = \{e_1, \dots, e_n\}$ its edge set. We always assume that G is simple, i.e., G has no loop and no multiple edge. Let $\mathbf{e}_1, \dots, \mathbf{e}_d$ denote the i th unit coordinate vectors of $\mathbb{R}^d$. We associate each edge $e = \{i, j\} \in E(G)$ with the vector $\rho(e) = \mathbf{e}_i + \mathbf{e}_j \in \mathbb{R}^d$. The *edge polytope* is the convex polytope $\mathcal{P}_G \subset \mathbb{R}^d$ which is the convex hull of the finite set $\{\rho(e_1), \dots, \rho(e_n)\}$. Let $K[\mathbf{t}] = K[t_1, \dots, t_d]$ be the polynomial ring in d variables over a field K . We associate each edge $e = \{i, j\} \in E(G)$ with the quadratic monomial $\mathbf{t}^e = t_i t_j \in K[\mathbf{t}]$. The *edge ring* is the affine semigroup ring $K[G] = K[\mathbf{t}^{e_1}, \dots, \mathbf{t}^{e_n}]$.

Let, in general, $\mathcal{P} \subset \mathbb{R}^d$ be an integral convex polytope, i.e., a convex polytope all of whose vertices have integer coordinates, which lies on a hyperplane $\mathcal{H} \subset \mathbb{R}^d$ with $\mathbf{0} \notin \mathcal{H}$, where $\mathbf{0}$ is the origin of $\mathbb{R}^d$. We assume that $\mathcal{P} \subset \mathbb{R}_{\geq 0}^d$, where $\mathbb{R}_{\geq 0}$ is the set of nonnegative real numbers. Then for each integer point $\mathbf{a} = (a_1, \dots, a_d)$ belonging to $\mathcal{P}$, we associate the monomial $\mathbf{t}^{\mathbf{a}} = t_1^{a_1} \cdots t_d^{a_d} \in K[\mathbf{t}]$. The *toric ring* of

2010 Mathematics Subject Classification: Primary 52B20; Secondary 13H10, 14M25.

Key words and phrases: finite graph, edge ring, edge polytope, Serre's condition (R_1) .

The first author is supported by the JST CREST “Harmony of Gröbner Bases and the Modern Industrial Society.”

This research was performed while the second author was staying at Department of Pure and Applied Mathematics, Osaka University, November 2011 – April 2012, supported by the DAAD.

$\mathcal{P}$ is the affine semigroup ring $K[\mathcal{P}] = K[\{\mathbf{t}^{\mathbf{a}} : \mathbf{a} \in \mathcal{P} \cap \mathbb{Z}^d\}]$. Thus in particular the edge ring $K[G]$ of a finite connected graph G is the toric ring of the edge polytope $\mathcal{P}_G$ of G .

We say that an integral convex polytope $\mathcal{P}$ is *normal* if its toric ring $K[\mathcal{P}]$ is normal. It is shown [3] and [6] that the edge ring of a finite connected graph is normal if and only if G satisfies the odd cycle condition [3, p. 410]. Recall that a toric ring is normal if and only if it satisfies Serre's condition (R_1) and (S_2) and that every normal toric ring is Cohen–Macaulay. Thus in particular a toric ring satisfying Serre's condition (R_1) is normal if and only if it is Cohen–Macaulay. In the present paper the problem when a given edge ring satisfies Serre's condition (R_1) is investigated.

2. WHEN DOES AN EDGE RING SATISFY SERRE'S CONDITION (R_1) ?

Let G be a finite connected graph on the vertex set $[d] = \{1, \dots, d\}$. If G is bipartite, then $K[G]$ is normal and satisfies Serre's condition (R_1) . Thus in what follows we assume that G is nonbipartite, i.e., G possesses at least one odd cycle.

If T is a nonempty subset of $[d]$, then the induced subgraph of G on T is denoted by G_T . A vertex $i \in [d]$ is called *regular* if every connected component of $G_{[d] \setminus i}$ has at least one odd cycle. A nonempty subset T of $[d]$ is called *independent* if $\{i, j\} \notin E(G)$ for all $i, j \in T$ with $i \neq j$. If $\emptyset \neq T \subset [d]$ is independent and if $N(G; T)$ is the set of vertices $j \in [d]$ with $\{i, j\} \in E(G)$ for some $i \in T$, then the *bipartite graph induced by T* is defined to be the bipartite graph having the vertex set $T \cup N(G; T)$ and consisting of all edges $\{i, j\} \in E(G)$ with $i \in T$ and $j \in N(G; T)$. Finally, we say that a nonempty subset $T \subset [d]$ is *fundamental* if

- T is independent;
- the bipartite graph induced by T is connected;
- either $T \cup N(G; T) = [d]$ or every connected component of the induced subgraph $G_{[d] \setminus (T \cup N(G; T))}$ has at least one odd cycle.

We are now in the position to state our criterion for an edge ring to satisfy Serre's condition (R_1) .

Theorem 2.1. *Let G be a finite connected graph on $[d]$ and suppose that G is nonbipartite. Then the edge ring $K[G]$ of G satisfies Serre's condition (R_1) if and only if the following conditions are satisfied:*

- (i) *For every regular vertex $i \in [d]$, the induced subgraph $G_{[d] \setminus i}$ is connected;*
- (ii) *For every fundamental set $T \subset [d]$, one has either $T \cup N(G; T) = [d]$ or the induced subgraph $G_{[d] \setminus (T \cup N(G; T))}$ is connected.*

Example 2.2. Let G be the finite connected graph on $\{1, \dots, 8\}$ with the edges 12, 23, 13, 45, 56, 46, 17, 47, 18 and 48, where each ij stand for the edge $\{i, j\}$. Since the graph G clearly violates the odd cycle condition, the edge ring $K[G]$ cannot be normal. The only vertices whose removal makes G disconnected are 1 and 4, but both are not regular. If $T \subset [8]$ is a set such that $G_{[d] \setminus (T \cup N(G; T))}$ is disconnected, then either 1 or 4 are contained in $T \cup N(G; T)$. But then $G_{[d] \setminus (T \cup N(G; T))}$ has only one odd cycle left, so T cannot be fundamental. Hence $K[G]$ satisfies Serre's condition

(R_1) . More generally, the same argument shows that the graphs G_{k+6} constructed in [5] satisfy (R_1) if and only if $k \geq 2$.

3. PROOF OF THEOREM 2.1

First recall the description of the facets of $\mathcal{P}_G$. To every regular vertex i we associate the linear form $\sigma_i : \mathbb{R}^d \rightarrow \mathbb{R}$ which projects onto the i -th component. Moreover, we set $\mathcal{H}_i = \{x \in \mathbb{R}^d : \sigma_i(x) = 0\}$ and $\mathcal{F}_i = \mathcal{P}_G \cap \mathcal{H}_i$. Similarly, to every fundamental set T we associate the linear form

$$\sigma_T : \mathbb{R}^d \ni (x_1, \dots, x_d) \mapsto \sum_{j \in N(G;T)} x_j - \sum_{i \in T} x_i$$

and we set $\mathcal{H}_T = \{x \in \mathbb{R}^d : \sigma_T(x) = 0\}$ and $\mathcal{F}_T = \mathcal{P}_G \cap \mathcal{H}_T$.

Lemma 3.1 ([3]). *The facets of $\mathcal{P}_G$ are exactly the sets $\mathcal{F}_i$ and $\mathcal{F}_T$ for all regular vertices i and all fundamental sets T .*

A combinatorial condition for a semigroup ring to satisfy Serre's condition (R_1) is explicitly stated in [1, Exercises 4.15, 4.16].

Proposition 3.2. *Let M be an affine monoid, K a field and $K[M]$ its semigroup ring. Then $K[M]$ satisfies Serre's condition (R_1) if and only if every facet $\mathcal{F}$ of M satisfies the following two conditions:*

- (i) *There exists $x \in M$ such that $\sigma_{\mathcal{F}}(x) = 1$, where $\sigma_{\mathcal{F}}$ is a support form of $\mathcal{F}$ taking integer values on $\text{gp}(M)$.*
- (ii) *$\text{gp}(M \cap \mathcal{F}) = \text{gp}(M) \cap \mathcal{H}$, where $\mathcal{H}$ is the supporting hyperplane of $\mathcal{F}$;*

where $\text{gp}(M)$ denotes the additive group generated by M .

We apply Proposition 3.2 to the affine monoid

$$M_G = \mathbb{N}(\mathcal{P}_G \cap \mathbb{Z}^d)$$

generated by the integer points in $\mathcal{P}_G$. Note that the support hyperplanes $\mathcal{H}_i$ and $\mathcal{H}_T$ of $\mathcal{P}_G$ are also the support hyperplanes of M_G . We start proving Theorem 2.1 by the following

Lemma 3.3. *The facets of M_G satisfy the first condition of Proposition 3.2.*

Proof. First, let $i \in [d]$ be a regular vertex. Since G is connected, there exists an edge $e = \{i, j\} \in E(G)$ to another vertex j . Then $\sigma_i(\rho(e)) = 1$.

Second, let $T \subset [d]$ be a fundamental set. If $T \cup N(G; T) \subsetneq [d]$, then there exists an edge $e = \{i, j\} \in E(G)$ such that $i \in N(G; T)$ and $j \in [d] \setminus (T \cup N(G; T))$. This edge satisfies $\sigma_T(\rho(e)) = 1$. If instead $T \cup N(G; T) = [d]$, then every edge of G has either both endpoints in $N(G; T)$, or one in $N(G; T)$ and one in T . Hence $\sigma_T(e) \in \{0, 2\}$ for every edge e of G . It then follows that $\frac{1}{2}\sigma_T$ satisfies the condition of Proposition 3.2. $\square$

To check the second condition of Proposition 3.2 we need to compute the lattice generated by M_G . The following Lemma 3.4 appears in [3, p. 426] without an explicit proof. However, for the sake of completeness, we give its detailed proof.

Lemma 3.4. *The lattice $gp(M_G)$ is the set of all integer vectors in $\mathbb{Z}^d$ with an even coordinate sum.*

Proof. Since every generator of M_G has an even coordinate sum, it follows that the lattice $gp(M_G)$ is contained in the set of all integer vectors in $\mathbb{Z}^d$ with an even coordinate sum.

To prove the converse, assume the edges $e_1, \dots, e_\ell$ form an odd cycle of G and let i be the common vertex of e_1 and e_ℓ . Then

$$2\mathbf{e}_i = \sum_{j=1}^{\ell} (-1)^{j+1} \rho(e_j) \in gp(M_G).$$

Now consider a spanning tree G' of G . The set $\{\rho(e) \mid e \in E(G')\}$ together with $2\mathbf{e}_i$ forms a $\mathbb{Z}$ -basis for the space of all integer vectors in $\mathbb{Z}^d$ with an even coordinate sum. $\square$

Now, we can prove two propositions, which complete our proof of Theorem 2.1.

Proposition 3.5. *Let $i \in [d]$ be a regular vertex of G . Then $\mathcal{F}_i$ satisfies the second condition of Proposition 3.2 if and only if $G_{[d] \setminus i}$ is connected.*

Proof. We denote the connected components of $G_{[d] \setminus i}$ with G'_j . Then it is easy to see that $M_{G_{[d] \setminus i}} = \bigoplus M_{G'_j}$ and hence $gp(M_G \cap \mathcal{F}_i) = gp(M_{G_{[d] \setminus i}}) = \bigoplus gp(M_{G'_j})$. Since every G'_j is connected and contains an odd cycle, we can use Lemma 3.4 to describe $gp(M_{G'_j})$. If $G_{[d] \setminus i}$ is connected, then $gp(G_{[d] \setminus i})$ and $gp(M_G) \cap \mathcal{H}_i$ are both the set of integer vectors in $\mathbb{Z}^d$ with even coordinate sum and i th coordinate equal to zero, thus these sets coincide.

We consider the case that $G_{[d] \setminus i}$ has at least two different connected components G'_1, G'_2 . Then we can choose a vector $x \in \mathbb{Z}^d$ such that (i) its coordinate sum is even, (ii) $\sigma_i(x) = 0$, and (iii) the restricted coordinate sum over the vertices in G'_1 is odd. This x is contained in $gp(M_G) \cap \mathcal{H}_i$, but not in $gp(G_{[d] \setminus i})$, thus $\mathcal{F}_i$ violates the condition. $\square$

Proposition 3.6. *Let $T \subset [d]$ be a fundamental set of G . Then $\mathcal{F}_T$ satisfies the second condition of Proposition 3.2 if and only if one has either $T \cup N(G; T) = [d]$ or the induced subgraph $G_{[d] \setminus (T \cup N(G; T))}$ is connected.*

Proof. Again, we denote the connected components of $G_{[d] \setminus (T \cup N(G; T))}$ with G'_j . We claim that

$$gp(M_G \cap \mathcal{F}_T) = \bigoplus gp(M_{G'_j}) \oplus \{x \in \mathbb{Z}^d \mid \text{supp}(x) \subset T \cup N(G; T), \sigma_T(x) = 0\}.$$

Here, $\text{supp}(\cdot)$ denotes the support of a vector. The sum is direct, because the supports of the summands are disjoint. $M_G \cap \mathcal{F}_T$ (and thus $gp(M_G \cap \mathcal{F}_T)$) is generated by the set $\{\rho(e) \mid e \in E(G), \sigma_T(\rho(e)) = 0\}$. For an edge $e \in E(G)$, it holds that $\sigma_T(\rho(e)) = 0$ if and only if either both endpoints lie in $T \cup N(G; T)$ or both are not contained in this set. Thus, a set of generators of the left side of (3) is contained in the right side of the equation, and hence one inclusion follows. Furthermore $\bigoplus gp(M_{G'_j}) \subset gp(M_G \cap \mathcal{F}_T)$. Thus it remains to show that

$$\{x \in \mathbb{Z}^d \mid \text{supp}(x) \subset T \cup N(G; T), \sigma_T(x) = 0\} \subset gp(M_G \cap \mathcal{F}_T).$$

For this we consider a spanning tree of the induced bipartite graph on $T \cup N(G; T)$. Its edges form a $\mathbb{Z}$ -basis for the left set, hence it is contained in $gp(M_G \cap \mathcal{F}_T)$. Next, we note that

$$gp(M_G) \cap \mathcal{H}_T = \left\{ x \in \mathbb{Z}^d \mid \text{supp}(x) \cap (T \cup N(G; T)) = \emptyset, \sum x_i \text{ even} \right\} \\ \oplus \left\{ x \in \mathbb{Z}^d \mid \text{supp}(x) \subset (T \cup N(G; T)), \sigma_T(x) = 0 \right\} .$$

Now the reasoning is completely analogous to the proof of Proposition 3.5. $\square$

REFERENCES

- [1] W. Bruns and J. Gubeladze, “Polytopes, Rings and K-Theory,” Springer–Verlag, 2009.
- [2] W. Bruns and J. Herzog, “Cohen–Macaulay rings,” Second Ed., Cambridge Univ. Press, 1998.
- [3] H. Ohsugi and T. Hibi, Normal Polytopes Arising from Finite Graphs, *J. Algebra* **207** (1998), 409–426.
- [4] H. Ohsugi and T. Hibi, Toric ideals generated by quadratic binomials, *J. Algebra* **218** (1999), 509–527.
- [5] T. Hibi, A. Higashitani, K. Kimura and A.B. O’Keefe, Depth of edge rings arising from finite graphs, *Proc. Amer. Math. Soc.* **139** (2011), 3807–3813.
- [6] A. Simis, W. V. Vasconcelos and R. H. Villarreal, The integral closure of subrings associated to graphs, *J. Algebra* **199** (1998), 281–289.

DEPARTMENT OF PURE AND APPLIED MATHEMATICS, GRADUATE SCHOOL OF INFORMATION
SCIENCE AND TECHNOLOGY, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN
E-mail address: hibi@math.sci.osaka-u.ac.jp

FACHBEREICH MATHEMATIK UND INFORMATIK, PHILIPPS-UNIVERSITÄT MARBURG, 35032
MARBURG, GERMANY
E-mail address: katthaen@mathematik.uni-marburg.de